



1

---

---

---

---

---

---

---

---



2

---

---

---


---

---

---

---

---



3

---

---

---

---

---

---

---

---



4

---

---

---

---

---

---

---

---

Different food → high JH in Q larvae →  
inhibits "programmed cell death" →  
more ovarioles

5

---

---

---

---

---

---

---

---



6

---

---

---


---

---

---

---

---




---

---

---

---

---

---

---

---

7




---

---

---

---

---

---

---

---

8




---

---

---

---


---

---

---

---

9




---

---

---

---

---

---

---

---

---

---

10

**The same pheromone is important for mating with drones**

Context dependent: drones do not respond to the same pheromone while inside the hive. Only while flying in the `drone-congregation` area.

We do not know what is the switch

---

---

---

---

---

---

---

---

---

---

11

**Virgin Queen Mating Areas and Drone Congregation Areas (DCA's)**

by [Gerald M. Lopez](#)



Drones fly out from their colonies every afternoon (season and weather permitting) in a search for virgin queens. Studies from other parts of the world and the United States conducted in the 1960's-1970's discovered that drones congregate in specific aerial locations day after day and year after year. These DCA's are of various sizes but often about 100m in diameter and, in these areas, they will fly higher (up to 60m) than the 12-20m elevation they fly as they leave the spairy or fly between DCA's. Beginning in 1965, we used a modified radar with a rotating parabolic-dish to track the drones and to locate DCA's.



This picture shows the radar screen in which a large DCA was observed. First, at the top of the picture, from left to right, the numbers indicate that the year was 1969, on the 89th day. The next numbers indicate that it was 3:01 MST. Just below the time is the angular elevation of the radar beam above the horizontal. Inside the circles, the center is where the radar is located, the first circle is a radius of 463 meters and the outer circle has a radius of 706 meters. This is a large area, or actually a volume of space was being "viewed" every 2 seconds (dish rotation rate). All the dots inside the circles are drones; in this case drone honey bees. As you can see, many drones are congregated in a particular area (the drones in this view were between 40-50m above ground) about 100m wide and 200m long. Actually, the drones in this picture are seen as a "slice" through the middle of the DCA. Even more drones were seen at the 40 and 50 "slices" and the top of the DCA was seen in the 7 to "slice" topping out at 60m above ground. By moving the radar and making such observations, we found 28 such DCA's and documented the interconnecting flyways between them.

These studies are written in the paper "Honey bee drone flyways and congregation areas, radar observations", 1992 in *J. Kansas Entomological Society*, 65:223-230 written by G. M. Lopez, W. W. Wolf, and O. R. Taylor, Jr.

---

---

---

---

---

---

---

---

---

---

12



<http://www.pbs.org>

13

---

---

---


---

---

---

---

---



14

---

---

---

---

---

---

---


---

**The only fecal stuff that smells good...  
only by the virgin queen!**

Virgin queens produce a chemical (in her defecation) that repels workers:

Chemical name:

o-aminoacetophenone.



15

---

---

---

---

---

---

---

---



16

---

---

---

---

---

---

---

---

Whose egg is it? The queen's or the workers?

We cannot tell, but the workers can!

17

---

---

---


---

---

---

---

---



18

---

---

---


---

---

---

---

---



19

---

---

---

---

---

---

---

---

---

---

**Worker policing:**

The removal of worker eggs by other workers (usually workers of another subfamily).

Policing requires that

- 1). Workers have the ability to distinguish queen eggs from worker eggs.
- 2). Workers do remove worker laid eggs under natural conditions.

20

---

---

---

---

---


---

---

---

---

---



21

---

---

---

---

---


---

---

---

---

---



22

---

---

---

---

---

---

---

---

---

---

---

---

Another queen tidbit

Unmated queens, like workers will produce only drones (no sperm to fertilize eggs)

However, *Apis mellifera capensis* (cape bee), unmated queens or workers can produce female offsprings at a high rate 60%ish. By fusing the polar body back to the original cell during meiosis.

Our bees will have about 1% for unmated virgin queens. But usually only for the eggs from the first few days.

23

---

---

---

---

---

---

---

---

---

---

---

---

Queen mating statistics

Queens left their nucs on  $2.20 \pm 0.98$  flying days (min. 1; max. 5),

Most of the nuptial flights (82.49%) taking place between 13:00 and 16:00 h (Figure 2).

The earliest and latest departure time: 11:50 and 17:38 h, respectively.

Number of recorded flights per queen was  $5.04 \pm 3.11$  (min. 1; max. 16), with a maximum of seven flights of one queen on one day.

Daily number of nuptial flights per queen was  $2.30 \pm 1.35$ , with a mean duration of  $17.69 \pm 13.19$  min (min. 3.08; max. 57.07; Figure 3).

*Insects* 2014, 5(3), 513-527;  
doi:10.3390/insects5030513

24

---

---

---

---

---

---

---

---

---

---

---

---





25

---

---

---

---

---

---

---

---



26

---

---

---

---

---

---

---

---

Drone Biology

Drones have no father (haploid!)  
All their sperms are clones (100% identical)  
Most drones are haploid, but inbreeding results in diploid drones  
Diploid drones are sterile and eaten by workers before 4<sup>th</sup> instar  
Sexually mature around 10-12 days (queens: 5-6 days).  
Mating flights between 3-5 pm.  
Drones fly to "drone congregation areas" to mate  
They die during mating (endophalus explodes)  
They get kicked out of the colonies in the fall

27

---

---

---


---

---

---

---

---



28

---

---

---

---

---

---

---

---

---

---

Drone sperm numbers:

1. Varroa mites reduce sperm numbers
2. Apistan reduces sperm numbers (Rinderer et al., 1999)
3. Smaller drones (drones reared in workers cells) have less sperms

29

---

---

---

---

---

---

---

---

---

---

From Duay, DeJone and Engles (2002)

*Flight performance and sperm production affected by Varroa* 231

**Table 1.** Flight performance and sperm production of drones in relation to the level of pupal infestation by *Varroa destructor*. Mean values, m = median, SD = standard deviation, r = range, d = diminished in relation to the controls.

Degree of pupal infestation of the drones	Duration of the test flights	Number of spermatozoa
Unparasitized	N = 64, $\bar{x}$ = 6'48" m = 4'54" SD = 5'35" r = 0'09" - 27'27"	N = 68 $\bar{x}$ = 7,540,441 m = 7,475,000 SD = 2,812,780 r = $2.5 \times 10^6$ - $12.8 \times 10^6$
One female mite per brood cell	N = 37 $\bar{x}$ = 6'55" m = 5'02" SD = 6'40" r = 0'15" - 22'15"	N = 53 $\bar{x}$ = 5,734,623 m = 4,200,000 SD = 3,574,404 r = $1 \times 10^6$ - $13.5 \times 10^6$ d = -24%
Two female mites per brood cell	N = 16 $\bar{x}$ = 2'16" m = 2'27" SD = 1'40" r = 0'09" - 6'01"	N = 31 $\bar{x}$ = 4,192,258 m = 3,550,000 SD = 2,506,754 r = $1 \times 10^6$ - $9.5 \times 10^6$ d = -45%

30

---

---

---

---

---

---


---

---

---

---

Smaller drones have less sperms (Schluns et al 2003)



31

---

---

---

---

---

---

---

---

### Controlling genetics in breeding:

1. Isolated mating yard (20 km)
2. Lots of drones (flooding, saturation)
3. Instrumental insemination
  
4. Late hours mating

32

---

---

---

---

---

---

---

---

### Late hour mating method

1. Confine both drones and queens with excluder material
2. Both must be sexually mature (drones: 10 days +, queen: 6 +)
3. Release both at the same time at 6 pm during summer.
4. Close again at 10 pm.
5. Repeat for 6 days.

### Post-test

33

---

---

---

---

---

---

---

---



34

---

---

---

---

---

---

---